

Primordial Black Holes and r-Process Nucleosynthesis

Volodymyr Takhistov (UCLA)



WIN-2017 @ UC-Irvine
(6.20.2017)

Based on: George Fuller, Alex Kusenko, VT [arXiv:1704.01129] (*accepted to PRL*)

- Dark matter (DM) nature unknown beyond gravitational interactions

Motivation: PBH DM

- Dark matter (DM) nature unknown beyond gravitational interactions
- Primordial black holes (PBH) could form in early Universe (proposed 50 yrs. ago)
 - **could be DM !**

[Zel'dovich, Novikov, 67;
Hawking, 71;
Carr, Hawking, 74]

Motivation: PBH DM

- Dark matter (DM) nature unknown beyond gravitational interactions
 - Primordial black holes (PBH) could form in early Universe (proposed 50 yrs. ago)
 - **could be DM !**
- [Zel'dovich, Novikov, 67;
Hawking, 71;
Carr, Hawking, 74]
- [Bird, Kamionkowski+, 16]
- Renewed interest: GW detection (PBH?), novel production mechanisms/signatures, no hints of popular DM particle candidates (e.g. WIMPs)

Motivation: PBH DM

- Dark matter (DM) nature unknown beyond gravitational interactions
- Primordial black holes (PBH) could form in early Universe (proposed 50 yrs. ago)
 - **could be DM !**

[Zel'dovich, Novikov, 67;
Hawking, 71;
Carr, Hawking, 74]
- Renewed interest: GW detection (PBH?), novel production mechanisms/signatures, no hints of popular DM particle candidates (e.g. WIMPs)

[Bird, Kamionkowski+, 16]
- PBH appear in many BSM scenarios and strictly, don't require non-SM physics
 - **plausible that regardless of DM origin, some in PBH !**

Motivation: PBH formation

- PBH formation: density contrast $\frac{\delta\rho}{\rho} \sim \mathcal{O}(1)$ within horizon \rightarrow collapse to BH

... improbable without new physics

see reviews

[Carr, Kuhnel, Sandstad, 17;
Khlopov, 10]

Motivation: PBH formation

- PBH formation: density contrast $\frac{\delta\rho}{\rho} \sim \mathcal{O}(1)$ within horizon \rightarrow collapse to BH
... *improbable without new physics*

see reviews
[Carr, Kuhnel, Sandstad, 17;
Khlopov, 10]
- Many early Universe production mechanisms
(e.g. scalar fields, baryogenesis, inflation, phase transitions, topological defects)

Motivation: PBH formation

- PBH formation: density contrast $\frac{\delta\rho}{\rho} \sim \mathcal{O}(1)$ within horizon \rightarrow collapse to BH
... *improbable without new physics*

see reviews
[Carr, Kuhnel, Sandstad, 17;
Khlopov, 10]
- Many early Universe production mechanisms
(e.g. scalar fields, baryogenesis, inflation, phase transitions, topological defects)
- Can estimate BH mass from formation time: $M_{\text{BH}} \sim 10^{15} \left(\frac{t}{10^{-23} \text{ s}} \right) g$

Motivation: PBH formation

- PBH formation: density contrast $\frac{\delta\rho}{\rho} \sim \mathcal{O}(1)$ within horizon \rightarrow collapse to BH
... improbable without new physics

see reviews
[Carr, Kuhnel, Sandstad, 17;
Khlopov, 10]
- Many early Universe production mechanisms
(e.g. scalar fields, baryogenesis, inflation, phase transitions, topological defects)
- Can estimate BH mass from formation time: $M_{\text{BH}} \sim 10^{15} \left(\frac{t}{10^{-23} \text{ s}} \right) \text{ g}$
- Thus, PBHs can span vast mass range (with mass spectrum):



General Setup

- If PBH form DM
 - many in DM-rich environments (e.g. Galactic Center)
- GC contains highest SN/star-formation rate
 - many neutron stars (NS), typically spinning (pulsars)


General Setup

- If PBH form DM
 - many in DM-rich environments (e.g. Galactic Center)
- GC contains highest SN/star-formation rate
 - many neutron stars (NS), typically spinning (pulsars)
- Thus, NS-PBH interactions ($\sim \rho_{\text{DM}} \times \rho_{\text{NS}}$) could be rather generic ...

General Setup

- If PBH form DM
 - many in DM-rich environments (e.g. Galactic Center)
- GC contains highest SN/star-formation rate
 - many neutron stars (NS), typically spinning (pulsars)
- Thus, NS-PBH interactions ($\sim \rho_{\text{DM}} \times \rho_{\text{NS}}$) could be rather generic ...
 - what are the astrophysical consequences?

Neutron Star Formation




Millisecond Pulsars

- Anticipating role of angular momentum in the amount of ejected material,
focus on pulsars with fastest rotation → millisecond pulsars (MSP)

Millisecond Pulsars

- Anticipating role of angular momentum in the amount of ejected material, focus on pulsars with fastest rotation → millisecond pulsars (MSP)
- Around 10-50% pulsars become MSP [Lorimer,13]
→ binary pulsar accretes matter from companion, spun-up and “recycled” → MSP

Image: NASA/Dana Berry



Millisecond Pulsars

- Anticipating role of angular momentum in the amount of ejected material, focus on pulsars with fastest rotation → millisecond pulsars (MSP)
- Around 10-50% pulsars become MSP [Lorimer,13]
→ binary pulsar accretes matter from companion, spun-up and “recycled” → MSP
- Population vs. rotation period: [Cordes,Chernoff,97; Lorimer,13]





Image: NASA/Dana Berry



NS-PBH Capture

- Case A: PBH captured during star formation → unlikely [Capela,Pshirkov,Tinyakov,13-14]

NS-PBH Capture


- ~~Case A: PBH captured during star formation - unlikely~~ [Capela, Pshirkov, Tinyakov, 13-14]
- Case B: PBH captured during NS lifetime

NS-PBH Capture

- ~~Case A: PBH captured during star formation~~ ~~unlikely~~ [Capela, Pshirkov, Tinyakov, 13-14]
- Case B: PBH captured during NS lifetime

Stage 1: gravitational capture

→ PBH approaches and passes through NS



NS-PBH Capture


- ~~Case A: PBH captured during star formation~~ ~~unlikely~~

[Capela, Pshirkov, Tinyakov, 13-14]

- Case B: PBH captured during NS lifetime

Stage 1: gravitational capture

- PBH approaches and passes through NS
- loses energy by dynamical friction f_{dyn}




NS-PBH Capture

- ~~Case A: PBH captured during star formation - unlikely~~ [Capela, Pshirkov, Tinyakov, 13-14]
- Case B: PBH captured during NS lifetime

Stage 1: gravitational capture

- PBH approaches and passes through NS
- loses energy by dynamical friction f_{dyn}
- if $E_{\text{loss}} > \text{KE}_{\text{PBH}} \rightarrow \text{captured !}$



NS-PBH Capture

- ~~Case A: PBH captured during star formation~~ ~~unlikely~~ [Capela, Pshirkov, Tinyakov, 13-14]
- Case B: PBH captured during NS lifetime

Stage 1: gravitational capture

- PBH approaches and passes through NS
- loses energy by dynamical friction \mathbf{f}_{dyn}
- if $E_{\text{loss}} > \text{KE}_{\text{PBH}} \rightarrow \text{captured !}$

capture rate:
$$F = \left(\frac{\Omega_{\text{PBH}}}{\Omega_{\text{DM}}} \right) F_0[m_{\text{PBH}}, \rho_{\text{DM}}, \bar{v}, R_{\text{NS}}, R_s]$$


NS-PBH Capture

- ~~Case A: PBH captured during star formation - unlikely~~ [Capela, Pshirkov, Tinyakov, 13-14]
- Case B: PBH captured during NS lifetime

Stage 2: PBH in NS

→ captured PBH continues passing through NS, until it settles inside

settle time: $t_{\text{set}} \simeq 4.1 \times 10^4 \left(\frac{m_{\text{PBH}}}{10^{22} \text{g}} \right)^{-3/2} \text{yr}$




NS-PBH Capture

- ~~Case A: PBH captured during star formation - unlikely~~ [Capela, Pshirkov, Tinyakov, 13-14]
- Case B: PBH captured during NS lifetime

Stage 3: BH grows inside

→ PBH inside NS grows via Bondi spherical accretion, consuming the host star

consume time: $t_{\text{con}} \simeq 10^{-2} \left(\frac{10^{22} \text{g}}{m_{\text{PBH}}} \right) \text{yr}$



Pulsar Lifetime


- Pulsar lifetime: $\langle t_{\text{NS}} \rangle = 1/F + t_{\text{set}} + t_{\text{con}}$
- Find $O(1 - 10)\%$ of NS consumed in Galactic time
→ consistent with observed missing pulsars [Dexter,O'Leary,14]

Bonus: consistent with recently discovered young GC magnetar [Mori+,13; Kennea+,13]
→ shows unusual activity ... a hint of BH consumption ??

Growing BH in NS: angular momentum transfer

- MSP spinning near mass shedding limit → elongated spheroid (Roche lobe model)


[Shapiro, Teukolsky, 83]



Growing BH in NS: angular momentum transfer

- MSP spinning near mass shedding limit → elongated spheroid (Roche lobe model)


[Shapiro, Teukolsky, 83]



Growing BH in NS: angular momentum transfer

- MSP spinning near mass shedding limit → elongated spheroid (Roche lobe model)

[Shapiro, Teukolsky, 83]




Add BH → can analytically show (see paper) surface matter exceeds escape speed
→ **ejected mass !!**

Growing BH in NS: angular momentum transfer

- MSP spinning near mass shedding limit → elongated spheroid (Roche lobe model)

[Shapiro, Teukolsky, 83]




Add BH → can analytically show (see paper) surface matter exceeds escape speed
→ **ejected mass !!**

* Differential rotation can occur → calculated that viscosity and magnetic stresses eliminate


Growing BH in NS: ejected mass

- Ejected mass:




Growing BH in NS: ejected mass

- Ejected mass:



Growing BH in NS: ejected mass


- Ejected mass:



- Population averaged: $\langle M_{\text{ej}} \rangle \sim \mathcal{O}(0.2) M_{\odot}$

Growing BH in NS: ejected mass

- Ejected mass:



- Population averaged: $\langle M_{\text{ej}} \rangle \sim \mathcal{O}(0.2) M_{\odot}$
- Ejecta neutron rich \rightarrow site of r-process nucleosynthesis?

- (R)apid-process nucleosynthesis:

[long list (Meyer, Schramm, *others*)]

- dominant mechanism for heavy element production
- neutrons capture on seed nuclei
- very sensitive to environment

R-process

- (R)apid-process nucleosynthesis:

[long list (Meyer, Schramm, *others*)]

- dominant mechanism for heavy element production
- neutrons capture on seed nuclei
- very sensitive to environment




Image: Los Alamos,
Nuclear Data Group

- (R)apid-process nucleosynthesis: [long list (Meyer, Schramm, *others*)]
 - dominant mechanism for heavy element production
 - neutrons capture on seed nuclei
 - very sensitive to environment
- Leading production sites: SN, compact object mergers (COM)
 - ... each has problems

R-process

- (R)apid-process nucleosynthesis: [long list (Meyer,Schramm, *others*)]
 - dominant mechanism for heavy element production
 - neutrons capture on seed nuclei
 - very sensitive to environment
- Leading production sites: SN, compact object mergers (COM)
 - ... each has problems
- PBH-NS: [Kouvaris,Tinyakov,13]
 - BH provides only slight heating → cold ejecta
 - neutrino emission negligible → don't spoil r-process [Meyer,McLaughlin,Fuller,98]

R-process

- (R)apid-process nucleosynthesis: [long list (Meyer,Schramm, *others*)]
 - dominant mechanism for heavy element production
 - neutrons capture on seed nuclei
 - very sensitive to environment
- Leading production sites: SN, compact object mergers (COM)
 - ... each has problems
- PBH-NS: [Kouvaris,Tinyakov,13]
 - BH provides only slight heating → cold ejecta
 - neutrino emission negligible → don't spoil r-process [Meyer,McLaughlin,Fuller,98]

Find amount of r-process material $O(10)$ larger than COM, several orders vs. SN !!

R-process: abundance from PBH-NS

- R-process material in Galaxy: $\sim 10^4 M_{\odot}$
- Recent UFD observations show consistency with 1 rare r-process event [Ji+,16]
→ difficult for SN, might be COM ... *how about PBH?*


R-process: abundance from PBH-NS

- R-process material in Galaxy: $\sim 10^4 M_{\odot}$
- Recent UFD observations show consistency with 1 rare r-process event [Ji+,16]
→ difficult for SN, might be COM ... *how about PBH?*

Amazingly, can explain both simultaneously with PBH-NS !!

* *studied for WIMP DM*
[Bramante,Linden,14]

R-process: abundance from PBH-NS



Other Signatures

Kilonova

- R-process nuclei build up and β -decay, resulting in EM emission
→ faint infrared after-glow days after event !

Other Signatures

Kilonova

- R-process nuclei build up and β -decay, resulting in EM emission
→ faint infrared after-glow days after event !

511-keV line

- Ejecta heated by β -decay and fission, generate many equilibrium positrons
→ find rate/energy matches with 511-keV line from positron-electron annihilation !

Other Signatures

Kilonova

- R-process nuclei build up and β -decay, resulting in EM emission
→ faint infrared after-glow days after event !

511-keV line

- Ejecta heated by β -decay and fission, generate many equilibrium positrons
→ find rate/energy matches with 511-keV line from positron-electron annihilation !

Fast Radio Bursts (FRB)

- Large energy release stored in magnetic flux tubes, if only (1-10)% of energy converted to radio waves → non-repeating FRB !

Summary

- PBH favorable DM candidate, expect in many scenarios
- PBH-NS interactions generic
- Analytically studied the setup:
 - site favorable to r-process, can explain amount in Galaxy + UFDs
 - mechanisms for other signatures (e.g. FRBs, 511-keV line, kilonova)

Summary

- PBH favorable DM candidate, expect in many scenarios
- PBH-NS interactions generic
- Analytically studied the setup:
 - site favorable to r-process, can explain amount in Galaxy + UFDs
 - mechanisms for other signatures (e.g. FRBs, 511-keV line, kilonova)

→ **possibility to resolve major astronomy puzzles simultaneously !**

Summary

- PBH favorable DM candidate, expect in many scenarios
- PBH-NS interactions generic
- Analytically studied the setup:
 - site favorable to r-process, can explain amount in Galaxy + UFDs
 - mechanisms for other signatures (e.g. FRBs, 511-keV line, kilonova)

→ **possibility to resolve major astronomy puzzles simultaneously !**

- Search identification:
 - no neutrino emission → easy to distinguish vs. SN
 - no gravity waves → easy to distinguish vs. COM

Summary

- PBH favorable DM candidate, expect in many scenarios
- PBH-NS interactions generic
- Analytically studied the setup:
 - site favorable to r-process, can explain amount in Galaxy + UFDs
 - mechanisms for other signatures (e.g. FRBs, 511-keV line, kilonova)

→ **possibility to resolve major astronomy puzzles simultaneously !**

- Search identification:
 - no neutrino emission → easy to distinguish vs. SN
 - no gravity waves → easy to distinguish vs. COM

→ **tested in upcoming experiments !**

Thank You for Attention!